

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Major Limitations in Using Element Concentrations in Hair as Biomarkers of Exposure to Toxic and Essential Trace Elements in Children

Helena Skräder, Maria Kippler, Barbro Nermell, Fahmida Tofail, Michael Levi, Syed Moshfiquur Rahman, Rubhana Raqib, and Marie Vahter

Table of Contents

Table S1. Recommended values for hair reference material NCSZC81002b, obtained values and limits of detection (LOD; $\mu\text{g/kg}$) for multiple elements in hair.

Table S2. Recommended values for reference materials, obtained values and limits of detection (LOD; $\mu\text{g/L}$) for multiple elements in urine.

Table S3. Recommended values for two reference materials, obtained values and limits of detection (LOD; $\mu\text{g/L}$) for multiple elements in erythrocytes.

Table S4. Spearman correlations (p-value) between all analyzed elements in hair at 10 years of age (n=207).

Figure S1. Increase in trace element concentrations along the hair (from scalp outwards) in 19 girls. The colors represent times increase relative to the first two cm. Only arsenic and selenium decreased slightly. The 1st-2nd cm represent recent exposure, and the 7th-8th cm exposure 8-10 months previously.

Figure S2. Scatter plots and smoothed lowess lines for non-transformed arsenic (As) in hair, erythrocytes and urine (left), and log2-transformed concentrations (right).

Figure S3. Spearman's correlation coefficients between all measured elements in hair at 10 years of age (top) and loading plot of principal component one and two for the principal components analysis (bottom).

Table S1. Recommended values for hair reference material NCSZC81002b, obtained values and limits of detection (LOD; $\mu\text{g/kg}$) for multiple elements in hair.

Element	N	Recommended value (mean \pm SD)	Obtained value (mean \pm SD)	LOD
Mg (mg/kg)	97	248 \pm 14	258 \pm 22	0.042
Ca (mg/kg)	97	1537 \pm 68	1703 \pm 92	0.12
Mn	97	3800 \pm 390	3273 \pm 308	0.10
Fe (mg/kg)	97	160 \pm 16	153 \pm 25	0.005
Co	97	150 \pm 15	175 \pm 69	<0.001
Cu (mg/kg)	97	34 \pm 2.3	35 \pm 3.0	<0.001
Zn (mg/kg)	97	191 \pm 16	194 \pm 11	0.009
As	97	200 \pm 2.3	196 \pm 21	0.014
Se	27	590 \pm 40	593 \pm 40	0.022
Mo	97	1100 \pm 100	1020 \pm 121	0.16
Cd	97	72 \pm 10	55 \pm 6.7	0.008
Pb	97	3830 \pm 180	4377 \pm 792	0.14

Table S2. Recommended values for reference materials, obtained values and limits of detection (LOD; µg/L) for multiple elements in urine.

Element	N	Serionorm™ Trace Elements Urine; 1011645		Serionorm™ Trace Elements Urine; 1011644		LOD
		Recommended value (mean ± SD) / Range ^a	Obtained value (mean ± SD)	Recommended value (mean ± SD) / Range ^a	Obtained value (mean ± SD)	
Mn	15	11 ± 2.2 / 6.5-15.3	10 ± 0.29	0.73 ± 0.15 / 0.44-1.02	0.73 ± 0.050	0.004
Fe	15	14 ^b	17 ± 4.7	14 ^a	15 ± 2.9	0.27
Co	15	11 ± 2.1 / 6.4-14.8	9.8 ± 0.40	0.72 ± 0.15 / 0.43-1.01	0.67 ± 0.023	0.002
Cu	15	22 ^b	20 ± 1.1	31 ^b	26 ± 1.2	0.078
Zn	15	1338 ± 269 / 800-1876	1131 ± 63	334 ± 67 / 200-468	295 ± 15	0.25
As	15	184 ± 37 / 110-258	184 ± 4.5	79 ± 16 / 47-111	81 ± 1.5	0.042
Se	15	70 ± 14 / 41.9-98.3	74 ± 4.4	14 ± 2.8 / 8.3-19.5	15 ± 0.46	0.004
Mo	15	38 ^b	39 ± 2.2	37 ^b	40 ± 1.8	0.034
Cd	15	4.9 ± 0.20 / 4.4-5.4	5.0 ± 0.12	0.20 ± 0.040 / 0.13-0.27	0.15 ± 0.0060	<0.001
Pb	15	91 ± 18 / 54.1-127.3	86 ± 2.8	0.66 ± 0.13 / 0.39-0.93	0.54 ± 0.031	<0.001

^a Acceptable range (SERO's assessment)

^b Approximate value

Table S3. Recommended values for two reference materials, obtained values^a and limits of detection (LOD; µg/L) for multiple elements in erythrocytes.

		Seronorm TM Trace Elements Whole Blood L-1; LOT 1103128		Seronorm TM Trace Elements Whole Blood L-2; LOT 1103129			
Element	N	Recommended value (mean ± SD) / Range ^b	Obtained value (mean ± SD)	N	Recommended value (mean ± SD) / Range ^b	Obtained value (mean ± SD)	LOD
Mn	29	21 ± 4.2 / 12.3-29.1	20 ± 0.40	27	30 ± 6.0 / 17.9-41.9	30 ± 0.59	0.012
Fe (mg/kg)	29	331 ^c	287 ± 15	27	319 ^c	286 ± 15	0.31
Co	29	0.16 ± 0.030 / 0.10-0.22	0.16 ± 0.030	27	5.8 ± 1.2 / 3.5-8.1	5.5 ± 0.15	0.003
Cu	29	680 ± 140 / 410-950	593 ± 19	27	1330 ± 270/ 790-1870	1255 ± 39	0.047
Zn	29	4400 ± 200 /4000-4800	3999 ± 124	27	6500 ± 300/ 5800-7200	6252 ± 180	0.68
As	29	2.4 ± 0.5 / 1.4-3.4	2.3 ± 0.21	27	14 ± 2.9 / 8.5-20.1	15 ± 1.4	0.014
Se	29	59 ± 12 / 35-83	58 ± 1.5	27	112 ± 23 / 66-158	121 ± 2.5	0.006
Mo	29	0.94 ± 0.22 / 0.61-1.27	1.0 ± 0.072	27	6.4 ± 1.3 / 3.84-9.00	6.5 ± 0.29	0.002
Cd	29	0.36 ± 0.020 /0.32-0.40	0.31 ± 0.033	27	5.8 ± 0.20 7 / 5.4-6.2	5.8 ± 0.12	0.003
Pb	29	10 ± 2.1 / 6.0-14.4	10 ± 0.23	27	310 ± 62 / 186-434	316 ± 3.7	0.009

^a Obtained values (originally µg/kg) recalculated to µg/L assuming a blood density of 1.060 g/ml

^b Acceptable range (SERO's assessment)

^c Additional value

Table S4. Spearman correlations (p-value) between all analyzed elements in hair at 10 years of age (n=207).

	Ca	Mg	Fe	Co	Cu	Zn	As	Se	Mo	Cd	Pb
Mg	0.76 (<0.001)	0.43 (<0.001)	0.46 (<0.001)	0.62 (<0.001)	0.098 (0.16)	0.28 (<0.001)	0.073 (0.30)	0.069 (0.32)	0.083 (0.24)	0.066 (0.34)	-0.049 (0.48)
Ca		0.55 (<0.001)	0.31 (<0.001)	0.56 (<0.001)	0.062 (0.37)	0.27 (<0.001)	0.0015 (0.98)	0.062 (0.37)	0.12 (0.081)	0.066 (0.35)	-0.099 (0.16)
Mn			0.23 (0.0011)	0.54 (<0.001)	0.15 (0.030)	-0.062 (0.37)	0.056 (0.42)	-0.12 (0.092)	0.23 (0.0010)	0.39 (<0.001)	0.40 (<0.001)
Fe				0.80 (<0.001)	-0.034 (0.62)	0.0020 (0.98)	0.13 (0.053)	0.18 (0.0085)	0.20 (0.0038)	0.029 (0.68)	0.053 (0.45)
Co					0.080 (0.25)	0.079 (0.26)	0.027 (0.70)	0.18 (0.0093)	0.24 (<0.001)	0.16 (0.020)	0.15 (0.035)
Cu						0.12 (0.096)	-0.028 (0.69)	0.040 (0.57)	0.13 (0.057)	0.21 (0.0027)	0.25 (<0.001)
Zn							-0.064 (0.36)	0.23 (<0.001)	-0.065 (0.35)	-0.22 (0.0015)	-0.48 (<0.001)
As								-0.011 (0.88)	0.24 (<0.001)	0.072 (0.30)	0.082 (0.24)
Se									0.25 (<0.001)	0.029 (0.67)	-0.098 (0.16)
Mo										0.17 (0.014)	0.19 (0.0071)
Cd											0.58 (<0.001)

Element	Hair segments from scalp outwards			
	1 st -2 nd cm	3 rd -4 th cm	5 th -6 th cm	7 th -8 th cm
Magnesium				
Calcium				
Manganese				
Iron				
Cobalt				
Copper				
Zinc				
Arsenic				
Selenium				
Molybdenum				
Cadmium				
Lead				

0.5-0.99	
1.0 (reference)	
1.01-1.5	
1.51-2.0	
2.0-2.5	
2.51-3.0	
3.01-3.5	
3.51-4.0	
4.01-4.5	
4.51-5.0	

Figure S1. Increase in trace element concentrations along the hair (from scalp outwards) in 19 girls. The colors represent times increase relative to the first two cm. Only arsenic and selenium decreased slightly. The 1st-2nd cm represent recent exposure, and the 7th-8th cm exposure 8-10 months previously.







Figure S2. Scatter plots and smoothed lowess lines for non-transformed arsenic (As) in hair, erythrocytes and urine (left), and log2-transformed concentrations (right).




Figure S3. Spearman's correlation coefficients between all measured elements in hair at 10 years of age (top) and loading plot of principal component one and two for the principal components analysis (bottom).